



US Army Corps
of Engineers®

ERDC/EL TN-16-2
May 2016

Technology Transfer of Biopolymer Soil Amendment for Rapid Revegetation and Erosion Control at Fort A. P. Hill, Virginia

by Steven Larson, W. Andy Martin, Roy Wade, Richard Hudson,
and Catherine Nestler

PURPOSE: Sustainable land management at Army installations is a critical concern of the modern Army and the Army engineer. A unique soil additive consisting of a polysaccharide polymeric material, a natural product of plant/soil rhizobial microbial activity, was demonstrated to enhance site vegetation and control erosion. The effort was supported by the Environmental Security Technology Certification Program (ESTCP, Project ER-0920) of the Department of Defense. *Rhizobium tropici*, a catalogued symbiotic nodulator of leguminous plants (Martinez-Romero et al. 1991), is also known for its prolific production of a gel-like, extracellular polymeric substance (EPS), a biopolymer (Gil-Serrano et al. 1990). The natural functions of the EPS in the rhizosphere include surface adhesion, self-adhesion of cells into biofilms, formation of protective barriers, water retention around roots, and nutrient accumulation (Laspidou and Rittmann 2002). The secretion of EPS by bacteria is recognized as a cohesive force in promoting surface erosion resistance in sediments (Droppo 2009, Gerbersdorf et al. 2008a, 2008b).

This biopolymeric material is produced commercially in dedicated bioreactors, separated from the growth media and derivatized in order to produce a non-reactive (non-crosslinking) material (Newman et al. 2010, Patent No. 7,824,569). Addition of *R. tropici* biopolymer to disturbed soil along with normal grass seeding activities has, through laboratory and field demonstrations, been shown to be an effective and low-cost method of reducing soil erosion and facility maintenance costs (Larson et al. 2012, Muller and Farr 2015). The biopolymer is non-toxic and — while long-lived — is ultimately biodegradable (MSDS). The case study described in this Technical Note (TN) evaluates application of this biopolymer to a highly disturbed and erodible soil.

BACKGROUND: Most training areas present soil erosion challenges from the production of dust and from reduced vegetation growth that results from heavy use (Figure 1). Maintenance of these areas is important for enhancing training land usage and availability.

At Fort A.P. Hill, any land disturbance greater than 2,500 ft² requires both an Erosion & Sediment (E&S) Control Plan and a Stormwater Pollution Prevention Plan (SWPPP). The field demonstration site, an airplane landing zone, is an area that has required extensive repair and maintenance over many years and has failed to meet the requirements outlined in the E&S Control Plan and the SWPPP.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE MAY 2016		2. REPORT TYPE		3. DATES COVERED 00-00-2016 to 00-00-2016	
4. TITLE AND SUBTITLE Technology Transfer of Biopolymer Soil Amendment for Rapid Revegetation and Erosion Control at Fort A. P. Hill, Virginia				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) U.S. Army Engineer Research and Development Center,,Vicksburg,,MS, 39180				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES The original document contains color images.					
14. ABSTRACT PURPOSE: Sustainable land management at Army installations is a critical concern of the modern Army and the Army engineer. A unique soil additive consisting of a polysaccharide polymeric material, a natural product of plant/soil rhizobial microbial activity, was demonstrated to enhance site vegetation and control erosion. The effort was supported by the Environmental Security Technology Certification Program (ESTCP, Project ER-0920) of the Department of Defense. Rhizobium tropici, a catalogued symbiotic nodulator of leguminous plants (Martinez-Romero et al. 1991), is also known for its prolific production of a gel-like, extracellular polymeric substance (EPS), a biopolymer (Gil-Serrano et al. 1990). The natural functions of the EPS in the rhizosphere include surface adhesion, self-adhesion of cells into biofilms, formation of protective barriers, water retention around roots, and nutrient accumulation (Laspidou and Rittmann 2002). The secretion of EPS by bacteria is recognized as a cohesive force in promoting surface erosion resistance in sediments (Droppo 2009, Gerbersdorf et al. 2008a, 2008b).					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES 9	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			



Figure 1. Examples of the maintenance challenges posed by training on military installations.

OBJECTIVE: The objective of the Fort A.P. Hill field study was to demonstrate the effectiveness and ease of biopolymer application. The biopolymer application can be completed using maintenance equipment normally found on Armed Forces installations and as part of routine installation maintenance in order to meet installation soil erosion control requirements.

SOIL CHARACTERIZATION: Bulk site soil samples were extracted and analyzed for metals using USEPA SW846 Method 3050. Metals analysis was performed by inductively coupled plasma mass spectrometry (ICP-MS). Soil pH was determined by electrode. The soil is characterized as a Sandy Loam. The metals analysis is shown in Table 1. Only one metal, aluminum (Al), shaded red, was present at a concentration well above the national average and above the range of concentration found across the conterminous United States (Shacklette and Boergen 1984). Aluminum is known to be toxic to plants at this concentration (Kochian 1995, Ma and Hiradate 2000, Ma et al. 2001; Ryan and Kochian 1993, and others). Soil pH was 4.73.

Table 1. Concentrations (mg/Kg) of select metals in the soil of the Fort A.P. Hill demonstration site as compared to national averages.			
Metal	Fort A.P. Hill (avg/stdev, mg/Kg)	U.S. avg* (mg/Kg)	U.S. range* (mg/Kg)
Aluminum (Al)	340,600/3,470	72,000	700 - >10,000
Sodium (Na)	37,127/2,920	12,000	<500 - 100,000
Cobalt (Co)	1.6/0.1	91	<3 - 70
Magnesium (Mg)	264/3	9,000	50 - >100,000
Calcium (Ca)	135/24	24,000	100 - 320,000
Chromium (Cr)	10.2/0.2	54	1 - 2,000
Manganese (Mn)	12/0.3	550	<2 - 7,000
Iron (Fe)	10,589/223	26,000	100 - >100,000
Nickel (Ni)	3/0.2	19	<5 - 700
Copper (Cu)	14/2	25	<1 - 700
Zinc (Zn)	21/6	60	<5 - 2,900
Lead (Pb)	5/1	19	<10 - 700

*Shacklette and Boergen (1984)

FIELD DEMONSTRATION: The purpose of the field demonstration at Fort A.P. Hill was three-fold:

- To control soil erosion from runoff water on the site
- To increase the germination rate and density of the vegetation (native grasses)
- To provide technology transfer to the Fort A.P. Hill DPW personnel with hands-on training in the application of the biopolymer.

The field demonstration site was a 3.8-acre area off Mosbey Rd. near the Airfield Landing area. This site, constructed in 2003, had never been seeded and little volunteer vegetation had been established. The site was divided into Area 1 and Area 2 (Figure 2); both consisted of a mostly sandy loam soil with 3-4-foot-deep ruts, small woody vegetation, some weeds, and little to no grass (Figure 3). The areas are adjacent to West Branch Mount Creek, which receives the runoff water.



Figure 2. Aerial view of the Fort A.P. Hill field demonstration.



Figure 3. Area 1 prior to site preparation.

The biopolymer, produced by NanoQuantics, Inc. (Wise, VA), a subsidiary of UXB International, was applied to both areas along with a custom grass seeding mix (700 lb). The seed mix was composed of

- Gulf annual ryegrass - 38.4%;
- VNS perennial ryegrass - 28.8%;
- Boreal red fescue - 29.51%; and
- other inert material - 3.29%.

The demonstration was begun on 26 May 2015. The Fort A.P. Hill Department of Public Works (DPW) determined that Area 1 and Area 2 required reconstruction prior to seeding. The site was prepared using construction equipment and trained operators available on the installation: bulldozers, a road grader, scraper, and tractors (Figure 4).



Figure 4. Earthwork using a bulldozer and road grader available, with trained operators, on-site.

Standard Operation and Maintenance (O&M) of grassed areas at Fort A.P. Hill requires soil fertilization and liming prior to seeding. After reconstruction of the site, biopolymer, fertilizer, lime, and grass seed were also applied to the soil. These activities also used equipment and trained operators available on the installation: a hydroseeder, a drill seeder, and a 3-point broadcast spreader.

1. The site was first disked with a tractor (Figure 5).
2. A tractor with a 3-point broadcast spreader spread 133, 40-pound bags (2.7 tons) of pellet lime over Areas 1 and 2 (Figure 5).
3. Next, 66 40-pound bags (1.3 ton) of 10-10-10 fertilizer was spread over Area 1 and 13, 50-pound bags (0.3 ton) of 20-0-20 fertilizer was spread over Area 2.

Note: These operations may not be required on all installations.



Figure 5. Disking the soil and broadcast application of lime and fertilizer using equipment and trained operators available on-site.

A cultipacker was used to place the seeds in the soil. A broadcast spreader also works well. The biopolymer was prepared by adding 500 gallons of water and 25 gallons of biopolymer — a 20:1 dilution ratio — in the hydroseeder, which homogenized the biopolymer solution (Figure 6).

Alternatively, the hydroseeder can be used to place seed and biopolymer at the same time.



Figure 6. Application of biopolymer and the seed mixture using a hydroseeder and broadcast spreader. Equipment and trained operators were available on-site.

At Fort A.P. Hill, hay bales were placed around the site perimeter to control potential erosion from heavy rains until grass was established (Figure 7).

This step is also optional if stormwater runoff is not a consideration.



Figure 7. Cultipacking and finishing the field using equipment and trained operators available on-site.

The results of the revegetation effort at Fort A.P. Hill are presented in Figure 8 and show successful germination and rapid increase in biomass. The effort to reduce soil erosion was proven successful within one week postplanting. During the first five days of June, immediately following completion of the seeding effort, Fort A.P. Hill received 2.9 in. of rainfall. Over 1.3 in. of this fell in one day (see Figure 8, left). The total rainfall for June was only 5.99 in. (National Climatic Data Center (NCDC), <http://www.ncdc.noaa.gov> accessed 9 September 2015). DPW personnel reported minimal erosion damage from the early rain events.¹




Figure 8. Rapid seed germination evident at seven days and significant biomass production at three weeks post-planting.

¹ Personal communication. 2015. John Parker, Environmental & Natural Resources Division, Fort A.P. Hill.

There is evidence from the metals analysis that establishment of vegetation on the site is slowed due to the acid pH of the soil combined with high concentrations of aluminum. Liming efforts promoted in the standard O&M guidelines of Fort A.P. Hill provide some protection for germinating seeds. However, aluminum is known to reduce the size of the taproot and the root branching topology (Delhaize and Ryan 1995, Ma et al. 2001, Serrano et al. 2011). When the pH increase produced from the lime has been expended, there is a decrease in plant biomass and eventual plant death. When the grass has died, the soil is no longer resistant to erosion by runoff water and wind. Evidence presented by Flis et al. (1993) suggests that the presence of cell surface components such as exopolysaccharides in the rhizosphere, like the *R. tropici* biopolymer, can lower the aluminum toxicity sufficiently to allow plant growth. The biopolymer soil amendment also promotes soil particle aggregation. This increases the resistance of the soil to erosion from surface water runoff (Larson et al. 2012).

SUMMARY AND CONCLUSIONS: The field demonstration at Fort A. P. Hill had the following objectives:

- To establish desired vegetation at a site previously unseeded
- To control soil erosion in an erodible soil on a disturbed site
- To demonstrate the ease of applying the biopolymer using maintenance equipment normally found on Armed Forces installations and as part of routine installation maintenance
- To improve vegetative growth on the site when biopolymer was used in conjunction with standard O&M procedures

These objectives were all successfully accomplished.

The protection of plants from aluminum toxicity in acid soils using exopolysaccharides should be further investigated.

ADDITIONAL INFORMATION: This TN was prepared by Dr. Steve Larson, Dr. W. Andy Martin, Roy Wade, and Richard Hudson of the Environmental Engineering Branch, Environmental Laboratory, U.S. Army Engineer Research and Development Center, and Catherine Nestler of Applied Research Associates, Inc. (ARA), Vicksburg, MS. Funding was provided by Headquarters Department of the Army (HQDA), Office of the Assistant Chief of Staff for Installation Management (OACSIM). The demonstration was supported by the Department of Public Works at Fort A.P. Hill, VA. For additional information, contact Dr. Steve Larson (601) 634-3431, steve.l.larson@erdc.dren.mil or Dr. Andy Martin (601) 634-3710 andy.martin@usace.army.mil.

This TN should be cited as follows:

Larson, S., W. A. Martin, R. Wade, R. Hudson, C. Nestler. 2016. *Technology transfer of biopolymer soil amendment for rapid revegetation and erosion control at Fort A.P. Hill, VA*. ERDC/EL TN-16-2. Vicksburg, MS: U.S. Army Engineer Research and Development Center.

REFERENCES

- Delhaize, E. and P. R. Ryan. 1995. Aluminum toxicity and tolerance in plants. *Plant Physiology* 107: 315-321.
- Droppo, I. C. 2009. Biofilm structure and bed stability of five contrasting freshwater sediments. *Marine and Freshwater Research* 60: 690-699.
- Flis, S. E., A. R. Glenn, M. J. Dilworth. 1993. Review: The interaction between aluminum and root nodule bacteria. *Soil Biology and Biochemistry* 25: 403-417.
- Gerbersdorf, S. A., T. Jancke, B. Westrich, D. M. Paterson. 2008a. Microbial stabilization of riverine sediments by extracellular polymeric substances. *Geobiology* 6: 57-69.
- Gerbersdorf, S. A., W. Manz, D. M. Paterson. 2008b. The engineering potential of natural benthic bacterial assemblages in terms of the erosion resistance of sediments. *FEMS Microbial Ecology* 66: 282-294.
- Gil-Serrano, A., A. S. del Junco, P. Tejero-Mateo. 1990. Structure of the extracellular polysaccharide secreted by *Rhizobium leguminosarum* var. *phaseoli* CIAT 899. *Carbohydrate Research* 204: 103-107.
- Kochian, L.V. 1995. Cellular mechanisms of aluminum toxicity and resistance in plants. *Annual Review of Plant Physiology and Plant Molecular Biology* 46: 237-260.
- Larson, S. L., J. K. Newman, C. S. Griggs, M. Beverly, and C. C. Nestler. 2012. *Biopolymers as an alternative to petroleum-based polymers for soil modification: Treatability studies*. ERDC TR-12-8. Vicksburg, MS: USACE Engineer Research and Development Center.
- Lapidou, C. S., and B. E. Rittmann. 2002. A unified theory for extracellular polymeric substances, soluble microbial products, and active and inert biomass. *Water Research* 36: 2711-2720.
- Ma, J. F. and S. Hiradate. 2000. Form of aluminum for uptake and translocation in buckwheat (*Fagopyrum esculentum* Moench). *Planta* 211: 355-360.
- Ma, J. F., P. R. Ryan, and E. Delhaize. 2001. Aluminum tolerance in plants and the complexing role of organic acids. *TRENDS in Plant Science* 6: 273-278.
- Martinez-Romero, E., L. Segovia, F. M. Mercante, A. A. Franco, P. Graham, M. A. Pardo. 1991. *Rhizobium tropici*, a novel species nodulating *Phaseolus vulgaris* L. beans and *Leucaena* sp. trees. *International Journal of Systemic Bacteriology* 41: 417-426.
- Muller, K. S. and J. V. Farr. 2015. *Cost benefit analysis of biopolymers, petroleum-based polymers and traditional soil stabilization methods*. White Paper 2015-1. West Point, NY: Department of Systems Engineering, USMA.
- Newman, J. K., D. B. Ringelberg, K. P. O'Connell, W. A. Martin, V. F. Medina, S. L. Larson. 2010. Patent 7,824,569. *Soluble Salt Produced from a Biopolymer and a Process for Producing the Salt*. Washington, DC: United States Patent and Trademark Office (USPTO).

- Ryan, P. R. and L. V. Kochian. 1993. Interaction between aluminum toxicity and calcium uptake at the root apex in near isogenic lines of wheat (*Triticum aestivum* L) differing in aluminum tolerance. *Plant Physiol.* 102: 957-982.
- Serrano, H. C., M. J. Pinto, M. A. Martins-Loução, C. Branquinho. 2011. How does an Al-hyperaccumulator plant respond to a natural field gradient of soil phytoavailable Al? *Science of the Total Environment* 409: 3749-3756.
- Shacklette, H. T. and J. G. Boerngen. 1984. *Element concentrations in soils and other surficial materials of the conterminous United States*. U.S. Geological Survey Professional Paper 1270. Washington, D.C.

NOTE: The contents of this technical note are not to be used for advertising, publication, or promotional purposes. Citation of trade names does not constitute an official endorsement or approval of the use of such products.